

Original Manuscript

# Comparison of the DNA damage response in BEAS-2B and A549 cells exposed to titanium dioxide nanoparticles

M. Biola-Clier<sup>1,2</sup>, D. Beal<sup>1,2</sup>, S. Caillat<sup>1,2</sup>, S. Libert, L. Armand<sup>1,2</sup>,  
N. Herlin-Boime<sup>3</sup>, S. Sauvaigo<sup>4</sup>, T. Douki<sup>1,2</sup> and M. Carriere<sup>1,2,\*</sup>

<sup>1</sup>Laboratoire Lésions des Acides Nucléiques, Université Grenoble Alpes, INAC, SYMMES, 38000 Grenoble, France,

<sup>2</sup>Laboratoire Lésions des Acides Nucléiques, CEA, INAC, SYMMES, 38000 Grenoble, France, <sup>3</sup>CEA, IRAMIS, NIMBE, LEDNA, 91191 Gif sur Yvette, France and <sup>4</sup>LXRepair, Parvis Louis Néel, 38000 Grenoble, France

\*To whom correspondence should be addressed: Tel: +33 43 878 0328; Fax: +33 43 878 5090. Email: [marie.carriere@cea.fr](mailto:marie.carriere@cea.fr)

Received 28 May 2016; Revised 9 October 2016; Editorial decision 11 October 2016; Accepted 18 October 2016.

## Abstract

For some decades production of titanium dioxide nanoparticle (TiO<sub>2</sub>-NP) has been increasing at a considerable rate; concerns as to the toxicity of these particles upon inhalation have been raised. Indeed, TiO<sub>2</sub>-NPs have been shown to induce significant genotoxicity and to adversely affect both major DNA repair mechanisms: base excision repair (BER) and nucleotide excision repair (NER). The aims of the present study were to (i) compare the genotoxicity of TiO<sub>2</sub>-NPs and their impact on DNA repair processes on A549 alveolar carcinoma and BEAS-2B normal bronchial lung cell lines and (ii) delve deeper into the mechanisms leading to these effects. To achieve these goals, TiO<sub>2</sub>-NPs effects on cytotoxicity, genotoxicity, DNA repair activity and DNA repair gene expression were investigated in both cell lines upon exposure to 1–100 µg/mL of anatase/rutile, 21 nm TiO<sub>2</sub>-NPs. Our results show that TiO<sub>2</sub>-NPs induce comparable cytotoxic and genotoxic responses in BEAS-2B and A549 cells. Functional response to DNA damage is observed in both cell lines and consists of an overall downregulation in DNA repair processes. When evaluating the relative importance of the two DNA repair pathways, we observed a lower impact on BER compared with NER activities, suggesting that repair of oxidatively generated DNA damage is still triggered in these cells. This response becomes measurable at 4 h of exposure in BEAS-2B but only after 48 h of exposure in A549 cells. The delayed response in A549 cells is due to an initial overall and intense downregulation of the genes encoding DNA repair proteins. This overall downregulation correlates with increased methylation of DNA repair gene promoters and downregulation of NRF2 and BRCA1, which may thus be considered as upstream regulators. These results strengthen the evidence that TiO<sub>2</sub>-NP induces indirect genotoxicity in lung cells, via modulation of DNA repair processes, and shed some light on the mechanisms behind this effect.

## Introduction

Titanium dioxide nanoparticles (TiO<sub>2</sub>-NPs) are among the top five NPs used in consumer products (1). Their annual production is estimated at 3800–7800 tonnes in the USA alone (2) and is continuously increasing (1). It therefore appears necessary to characterise

the impact of these NPs on the health of those who handle them and are likely to inhale them; inhalation was shown to be the main human exposure route in occupational settings. *In vitro* and *in vivo* experiments indicated that the main toxicity mechanisms induced by TiO<sub>2</sub>-NPs include induction of inflammation and oxidative stress, as well as genotoxicity (1,3–5). In rodents, exposure of the lungs to

high doses of TiO<sub>2</sub>-NPs led to tumour development (1). Based on this evidence, TiO<sub>2</sub> was classified by the International Agency for Research on Cancer (IARC) in Group 2B, possibly carcinogenic to humans (6).

Genotoxicity is often considered as a hallmark of NP-induced carcinogenesis. The effect can be direct, when the target of the genotoxic agent is DNA itself, or indirect, when the target is something other than DNA (7). *In vitro*, TiO<sub>2</sub>-NPs have been shown to accumulate in cells mainly by endocytosis and to distribute in intracytoplasmic compartments (8,9). Some reports also indicated their presence in the nucleus of cells (10–14). However, the proportion was minor compared with the amount in cytoplasmic compartments, making direct interaction between DNA and TiO<sub>2</sub>-NPs unlikely. The main mechanism reported in relation to TiO<sub>2</sub>-NP genotoxicity is the generation of reactive oxygen species (ROS) (15). Production of ROS is considered as direct genotoxicity (7), because these species may migrate to the nucleus where they can cause single- or double-strand breaks in DNA or cause the formation of oxidised bases such as 8-oxo-7,8-dihydro-2'-deoxyguanosine (8-oxo-dGuo). These phenomena were reported to occur in cells exposed to TiO<sub>2</sub>-NPs following both acute (12,16) and chronic exposure (17,18). TiO<sub>2</sub>-NPs have also been reported to induce indirect genotoxicity via mechanical interference with microtubules (7) which, among other outcomes, hinders mitotic progression (19). In addition, TiO<sub>2</sub>-NPs have been reported to induce epigenetic modification (7) to modulate the expression of DNA repair genes and proteins (15,20,21) and to reduce DNA repair ability through the nucleotide excision repair (NER) and base excision repair (BER) pathways (12).

In the human respiratory tract, based on their size, NPs are predicted to mainly deposit in the alveoli (22). Alveolar effects are frequently studied using the A549 cell line, derived from a human adenocarcinoma of the alveolar basal epithelium. However, the use of A549 cells in studies of DNA damage response has been criticised, because these cells are hypotriploid and they have an increased oxidative stress response (23) due to a mutation in the gene encoding Kelch-like family member 19 (KEAP1). *KEAP1* encodes a cytoplasmic protein that sequesters nuclear factor (erythroid-derived 2)-like 2 (NRF2) in the cytoplasmic compartment. When not tethered by KEAP1, NRF2 locates to the cell nucleus where it promotes the expression of genes encoding redox regulation proteins. Since the accepted paradigm for particle-induced genotoxicity is oxidative stress, the A549 response to NPs may therefore differ from that of other cell types.

The aims of the present study were to (i) compare the response of non-cancerous bronchial epithelial BEAS-2B cells with that of A549 epithelial alveolar cancer cells, particularly their DNA damage response when exposed to TiO<sub>2</sub>-NPs and (ii) delve deeper into the mechanisms of DNA damage response induced by TiO<sub>2</sub>-NPs. These two cell lines were exposed to 1–100 µg/mL NM105 TiO<sub>2</sub>-NPs with an average primary diameter of 21 nm [from the Joint Research Centre (JRC) library] for 4 h, 24 h or 48 h. The exposure medium consisted of cell culture medium without foetal bovine serum (FBS), as in our previous studies (9,12). Cell viability, DNA integrity, DNA repair ability and expression of DNA repair genes were evaluated in both cell lines. The mechanisms through which DNA repair genes were repressed was then further investigated in A549 cells by analysing the methylation state of DNA repair gene promoters and expression levels for three upstream regulators of DNA repair gene expression.

## Materials and methods

### Chemicals and NPs

Unless indicated otherwise, chemicals were purchased from Sigma-Aldrich and were >99% pure.

NM105 TiO<sub>2</sub>-NPs were obtained from the nanomaterial library at the JRC (Ispra, Italy). Their physico-chemical characteristics were determined to be as follows: spherical shape; primary particle size, 21 ± 9 nm; anatase:rutile ratio, 86:14%; specific surface area, 46 m<sup>2</sup>/g (<http://bookshop.europa.eu/fr/titanium-dioxide-nm-100-nm-101-nm-102-nm-103-nm-104-nm-105-pLBNA26637/>); and point of zero charge, 7.0 (9).

### NP suspension

NPs were dispersed in ultrapure water as previously described by ultrasonication for 30 min at 4°C on a Vibra Cell 75043 sonicator (Biolock Scientific) operated in pulse mode (1 s on/1 s off) at 28% amplitude (9), i.e. 16.7 W (24). After dispersion in water, the number-based hydrodynamic diameter was 70 ± 20 nm, with a polydispersity index (PDI) of 0.18 ± 0.04. Just before cell exposure, these water suspensions were diluted in FBS-free cell culture medium at a final concentration of 20 µg/mL, a concentration compatible with the Malvern zetasizer. In these conditions, the hydrodynamic diameter shifted to 720 ± 20 nm and the PDI to 0.50 ± 0.02, suggesting agglomeration. Forty-eight hours after dilution in exposure medium, the hydrodynamic diameter was >1, indicating that NPs were totally agglomerated.

### Cell culture and exposure conditions

BEAS-2B cells were purchased from ATCC (CRL-9609) and cultured in Dulbecco's Modified Eagle's medium (Gibco, 11960-044) containing 10% FBS, antibiotics and glutamine on collagen-coated Petri dishes and plates. Cells were exposed to 0–100 µg/mL of TiO<sub>2</sub>-NPs in FBS-free cell culture medium for 4, 24 or 48 h.

### Cytotoxicity assay

TiO<sub>2</sub>-NPs effect on cell viability was investigated on A549 and BEAS-2B cells using the 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyl tetrazolium bromide (MTT) assay. Cells were grown in 96-well plates, exposed to TiO<sub>2</sub>-NPs for 4, 24 or 48 h; exposure media were then replaced with 100 µL of MTT, and cells were incubated for a further 1 h at 37°C. The formazan crystals that formed inside cells were then dissolved in 100 µL of dimethyl sulphoxide. To limit possible NP interference with the assay, plates were centrifuged for 5 min at 3200 × g, and 50 µL of the supernatant was transferred to a new plate for absorbance reading, as previously described (24). Assays were independently reproduced three times, with six independent replicates for each experiment (i.e., *n* = 6).

### Genotoxicity assays

#### Comet assay

DNA strand breaks and alkali-labile sites induced by TiO<sub>2</sub>-NPs were visualised using the alkaline version of the comet assay (25). About 20 000 cells from each sample were deposited on each slide, and three slides were analysed per sample. Lysis was performed overnight at 4°C, and electrophoresis was carried out at 25 V (300 mA) for 30 min. Comets were stained using ethidium bromide and measured using comet assay IV (Perceptive Instrument, Bury St Edmunds, UK). This assay was independently reproduced three times,

with three independent biological replicates in each case, each of which was loaded on two different comet slides.

### 8-oxo-dGuo and 5-Me-dC quantification by HPLC-MS/MS

The levels of 8-oxo-dGuo and 5-methyl deoxycytosine (5-Me-dC) were measured by high-performance liquid chromatography–mass spectrometry (HPLC-MS/MS). DNA from frozen cell pellets (−80°C) was extracted using the DNeasy® Blood and Tissue kit (Qiagen) according to the manufacturer's procedure. An RNase treatment step was added after cell lysis, samples were incubated for 2 min with 400 µg of RNaseA. DNA was eluted in 100 µL of 0.1 mM deferoxamine to prevent spurious oxidation (26). DNA from 50 µL of sample was digested in two incubation steps. First, the pH was adjusted to pH 5.5 with the addition of 5 µL of buffer (100 mM succinic acid, 50 mM calcium chloride, 150 mM, 5 µM zinc sulphate, pH 5.5). Samples were then incubated for 2 h at 37°C in the presence of 2.5 U of nuclease P1, DNase II and 0.05 U of phosphodiesterase II. Then, Tris buffer (6 µL, 500 mM Tris, 1 mM ethylenediaminetetraacetic acid, pH 8) was added together with 2 U of alkaline phosphatase and 0.05 U of phosphodiesterase I. The sample was incubated for a further 2 h at 37°C. Hydrochloric acid (3 µL, 0.1 N) was added, and the tubes were centrifuged at 7000 × g for 5 min. The aqueous phase was collected and transferred into HPLC injection vials for HPLC-MS/MS analysis. Chromatographic separation was performed on a C18 reverse-phase Uptisphere ODB column (Interchim, Montluçon, France). Analytes were eluted using a gradient of methanol in 2 mM ammonium formate at a flow rate of 0.2 mL/min. Upstream of the mass spectrometer, the HPLC eluate was analysed in an ultraviolet detector set at 280 nm to quantify the amount of unmodified nucleosides. The retention time was around 29 min. MS/MS was carried out on an API 3000 mass spectrometer (SCIEX) in multiple reaction monitoring mode with positive electrospray ionisation, as previously described (12,26). The monitored fragmentation were  $m/z$  284 [M + H]<sup>+</sup> →  $m/z$  168 [M + H − 116]<sup>+</sup> for 8-oxo-dGuo and  $m/z$  242 →  $m/z$  126 for 5-Me-dC. Levels of 8-oxo-dGuo were expressed as a number of lesions per million normal bases. Levels of 5-Me-dC were expressed as % of lesions per dC. This assay was reproduced three times independently, with  $n = 3$  in each replicate.

### Analysis of DNA repair gene expression

DNA repair gene expression was assessed by reverse transcription quantitative polymerase chain reaction (RT-qPCR). Briefly, RNA was extracted and reverse-transcribed using the GenElute™ mammalian total RNA miniprep kit with the optional DNase treatment step and the SuperScript II Reverse Transcriptase kit (Life Technologies), according to the manufacturers' protocols. qPCR was performed on a MX3005P Multiplex Quantitative PCR thermocycler (Stratagene). Primer sequences are reported in supplementary Table 1, available at Mutagenesis Online. Three reference genes, GAPDH, S18 and CycloB were chosen for normalisation using BestKeeper (27). Relative quantification of gene expression and statistical analyses were performed with the REST 2009 software (28) based on the  $\Delta\Delta C_q$  method. A theoretical value of 2 was assumed for all PCR efficiencies. RT-qPCR was reproduced three times independently, with three independent biological replicates loaded twice on the qPCR plate.

### Multiplex array for DNA repair activity

The BER and NER abilities of BEAS-2B cells exposed to TiO<sub>2</sub>-NPs were assessed using a multiplexed excision/synthesis repair assay (LXRepair, Grenoble, France). Nuclear extracts were prepared as

described previously (29), and excision/repair reactions were run for 3 h at 30°C at a final protein concentration of 0.3 mg/mL on damaged plasmid microarrays (29). The reaction medium contained 1 mM adenosine triphosphate and 0.25 µM dCTP-Cy3 (GE Healthcare). The lesions assessed were photoproducts [cyclobutane pyrimidine dimers and (6–4) photoproducts (CPD-64)], 8-oxo-dGuo (8oxoG), alkylated bases (AlkB), abasic sites (AbaS) and pyrimidine glycols (Glycol). For each lesion, the total fluorescence incorporated into a lesion-containing plasmid was quantified using the Innoscan 710AL scanner (Innopsys, Toulouse, France) and normalised relative to the fluorescence incorporated into an undamaged plasmid. Each extract was tested twice, and the experiment was reproduced three times.

The GlycoSPOT assay (LXRepair, Grenoble, France) was used to quantify the excision activities of excision–resynthesis repair. In this assay, a multiplexed array of oligonucleotides (ODNs) containing specific lesions is used to monitor several glycosylases/AP endonuclease activities in cell extracts. The lesions comprised 8-oxoguanine paired with C (8oxoG-C), A paired with 8oxoguanine (A-8oxoG), ethenoadenine (EthA-T), thymine glycol (Tg-A), uracil (paired either with G or A (U-G and U-A respectively), hypoxanthine (Hx-T), abasic sites (THF-A). Cleavage of the lesions by the enzymes contained in the extracts released the fluorescence attached to the lesion-containing ODNs (30).

Repair reactions were allowed to proceed for 1 h at 37°C with 20 µg/mL of protein in 80 µL of excision buffer (10 mM HEPES/potassium hydroxide pH 7.8, 80 mM potassium chloride, 1 mM ethylene glycol-bis(β-aminoethyl ether)-N,N,N',N'-tetraacetic acid, 0.1 mM zinc chloride, 1 mM dithiothreitol and 0.5 mg/mL bovine serum albumin). After washing three times for 5 min at room temperature in phosphate-buffered saline containing 0.2 M sodium chloride and 0.1% Tween 20, the fluorescence emitted by spots was quantified using the Innoscan 710AL scanner. Each extract was run in duplicate. Data were normalised using the NormalizeIt software as described previously (29).

Percentage of excision of each lesion was calculated using the following formula:  $[100 \times (1\% \text{ of fluorescence of Lesion\_containing\_ODN} / \text{percentage of fluorescence of Lesion\_Free ODN})]$ . This calculation determines the percentage of residual fluorescence in the wells incubated with the extracts with respect to wells incubated in the buffer alone. Possible non-specific cleavage of the control ODN (Lesion-Free ODN) was also taken into account by the formula. These assays were performed once, with three independent biological replicates.

### Methylation in the promoters of DNA repair genes

DNA from frozen cell pellets was extracted using the DNeasy® Blood and Tissue kit (Qiagen), following the manufacturer's instructions with an extra RNase treatment step after cell lysis (2 min of incubation with 400 µg of RNase A, Sigma-Aldrich). DNA was eluted in nuclease-free water before digestion using the EpiTect® Methyl II DNA Restriction Kit (Qiagen) according to the manufacturer's recommendations. Samples were stored at −20°C. The methylation profile was analysed using the EpiTect® Methyl II Signature PCR Array Human DNA Repair kit (Qiagen) according to the manufacturer's protocol. qPCR was performed with MESA Blue qPCR Mastermix Plus for SYBR® Assay Low ROX (Eurogentec) on a MX3005P thermocycler (Stratagene). Data were analysed using the EpiTect Methyl II PCR Array Excel-based data analysis template, version 2.0 (Qiagen). This assay was independently reproduced three times, with  $n = 3$  on each replicate.

## Statistical analyses

For all experiments except qPCR, statistical significance was assessed with both non-parametric one-way analysis of variance on ranks (Kruskal–Wallis) and pairwise comparison by applying a Mann–Whitney *U* test using Statistica 8.0 software (Statsoft). Results were considered statistically significant when the *P* value was <0.05. The Relative Expression Software Tool (REST 2009) software was used to analyse qPCR data (28).

## Results

### Cytotoxicity

In both A549 and BEAS-2B cell lines, cytotoxicity induced by TiO<sub>2</sub>-NPs was determined using the MTT assay, which is based on the measurement of cellular NAD(P)H-dependent oxidoreductase enzymatic activity in viable cells. TiO<sub>2</sub>-NPs did not interfere with the MTT assay as previously shown (24). Cytotoxicity was observed only after 24 or 48 h exposure times, never at 4 h (Figure 1). However, more than 70% of the cells remained viable after 48 h of exposure, suggesting that TiO<sub>2</sub>-NP is moderately cytotoxic.


### Genotoxicity

Induction of strand breaks and alkali-labile sites by TiO<sub>2</sub>-NPs was assessed by performing the alkaline comet assay. Then 8-oxo-dGuo, which is the most frequently oxidised DNA base, was measured by HPLC-MS/MS. Cells were exposed to 10 or 100 µg/mL TiO<sub>2</sub>-NPs. At these concentrations, cell viability was >70%, as required for genotoxicity assays. These concentrations also limited possible interference of NPs with the assays. No DNA damage was induced at 10 µg/mL (Figure 2), in contrast, at 100 µg/mL, DNA damage was observed at all time points, in both cell lines and using both assays (Figure 2). Thus, comet tail intensity increased from 2–10% in control cells to 18–28% in cells exposed to 100 µg/mL TiO<sub>2</sub>-NPs (Figure 2A and B). Significantly higher levels of DNA damage were observed in A549 cells than in BEAS-2B cells at 4 h of exposure. However, at 48 h, damage was more extensive in BEAS-2B cells than in A549 cells, although the difference was modest. The number of 8-oxo-dGuo lesions ranged from 15 to 26 8-oxo-dGuo per million bases in exposed cells, while only three to eight 8-oxo-dGuo per million bases were detected in control cells (Figure 2C and D). This level of oxidation is within the acceptable range defined by the European standards committee on oxidative DNA damage (ESCODD) (31). The number of 8-oxo-dGuo lesions was moderately, but statistically significantly, higher in BEAS-2B than in A549 at 24 h of exposure (Figure 2C and D).


### DNA repair activity in cells exposed to TiO<sub>2</sub>-NPs

In A549 cells, we previously reported that 24 h of exposure to 100 µg/mL TiO<sub>2</sub>-NPs induces a slight increase in DNA repair activity through the BER and NER pathways, followed by a drastic reduction in DNA repair activity at 48 h post-exposure (12). This result was obtained using a microarray (ExSy-SPOT microarray) that measures repair efficiency for five DNA lesions, repaired either via the BER (8oxoG, AbaS, AlkB) or the NER (CPD-64, glycol) pathways (29). Here, the same experiment was carried out on BEAS-2B cells, exposed to 100 µg/mL TiO<sub>2</sub>-NPs for 4 h, 24 h or 48 h. At all three time points, exposure to TiO<sub>2</sub>-NPs induced a drastic reduction in cell's ability to repair the five lesions (not shown). Normalisation for the total fluorescence detected on the array was used to reveal whether a specific repair pathway played a more extensive role when compared with the overall DNA repair activity of the cell (supplementary Figure 1, available at Mutagenesis Online). As indicated above, AlkB, 8oxoG and AbaS are typically repaired by BER, while CPD-64 are repaired by NER. However, photoproducts and glycols (thymine and cytosine glycols) are transcription-blocking lesions and tend to cluster together in this assay. After normalisation of A549 (Figure 3A) and BEAS-2B data (Figure 3B), TiO<sub>2</sub>-NP exposure was found to induce a shift in the relative importance of two groups of repair activities. Thus, in control (unexposed) cells, CPD-64 and Glycols were more frequently repaired than AlkB, 8oxoG and AbaS, whereas the opposite effect was noted in TiO<sub>2</sub>-NP-exposed cells. A similar shift in repair activity was observed in A549 cells, but only after 48 h of treatment. These results suggest that although the DNA repair activity is decreased overall in BEAS-2B cells, the relative ability of the cell to repair oxidative lesions via the BER pathway is somehow preserved, while the NER pathway is more severely affected.

To further investigate the mechanisms by which TiO<sub>2</sub>-NP exposure modulates BER activity, we assessed their impact on the first step of this pathway, excision by glycosylases/AP endonucleases (32). To do this, we focused on A549 cells exposed for 24 h or 48 h to TiO<sub>2</sub>-NPs. The rationale for conducting this assay is that glycosylases—particularly OGG1, which is responsible for the excision of 8-oxo-dGuo—are susceptible to oxidative stress (33) and may thus be inactivated in TiO<sub>2</sub>-NP-exposed cells. Another multiplexed assay was used (GlycoSPOT), based on ODN cleavage (30). No significant impact was observed at 24 h of exposure to TiO<sub>2</sub>-NPs. However, after 48 h of exposure, the ability of A549 cells to excise adenine paired with 8-oxo-dGuo and EthA was decreased (Figure 4). Conversely, TiO<sub>2</sub>-NP exposure did not affect the cell's ability to excise uracil (U), 8-oxo-dGuo, thymine glycol (Tg), hypoxanthine (Hx) or abasic sites (THF-A) (Figure 4).



**Figure 1.** Cytotoxicity of TiO<sub>2</sub>-NPs towards A549 and BEAS-2B cells. A549 (A) and BEAS-2B (B) cell mortality was assessed using the MTT assay in cells exposed to 0–100 µg/mL TiO<sub>2</sub>-NPs for 24 or 48 h. The assays were independently reproduced three times, with *n* = 6 in each independent experiment.



**Figure 2.** TiO<sub>2</sub>-NP-induced genotoxicity in BEAS-2B and A549 cells. Genotoxicity was assessed using the alkaline comet assay (A, B) and by assaying 8-oxo-dGuo (C, D). A549 cells (A, C) and BEAS-2B cells (B, D) were exposed to 100 µg/mL of TiO<sub>2</sub>-NPs for 4, 24 or 48 h. Experiments were independently repeated three times with three biological replicates each time. Statistically significant results are indicated: \**P* < 0.05, exposed vs. control; #*P* < 0.05, A549 vs. BEAS-2B in the same exposure conditions.

### Modulation of the expression of genes encoding DNA repair proteins in TiO<sub>2</sub>-NP-exposed cells

To examine the mechanisms behind this deregulation of DNA repair activities, we measured the expression levels of 15 genes encoding DNA repair proteins from the BER or NER pathways by RT-qPCR. For these experiments, both cell lines were exposed to 1, 10 or 100 µg/mL TiO<sub>2</sub>-NPs. No modulation of gene expression was observed in cells exposed to 1 µg/mL TiO<sub>2</sub>-NPs and only one gene, *CSB*, was affected at 10 µg/mL (data not shown).

The most intense modulation of gene expression was observed in A549 cells exposed to 100 µg/mL TiO<sub>2</sub>-NPs for 4 h or 24 h. In these conditions, 12 of the 15 genes monitored were downregulated, of which 7 at 4 h (*XRCC1*, *PCNA*, *LIG3*, *XPC*, *DDDB1*, *DDB2* and *CSB*) and 11 at 24 h (*APE1*, *PARP1*, *POLB*, *PCNA*, *LIG3*, *bHR23B*, *XPC*, *DDDB1*, *DDB2*, *CSA* and *CSB*) (Table 1). The transcriptional response was much less intense at 48 h of exposure in this cell line—with downregulation of only one gene, *PARP1*—and in BEAS-2B cells at all time points. Indeed, in BEAS-2B cells exposed to 100 µg/mL TiO<sub>2</sub>-NPs, only 3 of the 15 genes were downregulated, 2 of them at 4 h and 24 h of exposure (*POLB* and *PCNA*) and 1 at 48 h (*bHR23B*) (Table 1).

Deregulated expression of three genes was common to both cell lines. These genes were, *POLB* at 4 h, *PCNA* at 4 h and 24 h and *bHR23B*, which was downregulated at 24 h in A549 and at 48 h in BEAS-2B (Table 1, boldface values). The fold change in gene expression for these genes was consistently lower in A549 cells than in BEAS-2B cells, indicating a more intense transcriptional regulation in A549.

Because DNA repair gene expression was more intensively modulated in A549 cells, we chose this cell line to further study the effects of TiO<sub>2</sub>-NP exposure on genes encoding other repair proteins involved in the BER and NER pathways, as well as genes linked to non-homologous end joining, homologous recombination (HR) and mismatch repair. Modulation of expression levels was primarily observed in cells exposed to 100 µg/mL TiO<sub>2</sub>-NPs


for 4 h or 24 h. A schematic representation of these DNA repair pathways is presented in Figure 5, together with the proteins for which gene expression was found to be deregulated. As shown in Figure 5 and in supplementary Table 2, available at Mutagenesis Online, the general trend was for downregulation, with 19 of the 29 genes tested (65%) downregulated, with fold changes ranging between 0.36 and 0.90 (supplementary Table 2, available at Mutagenesis Online). The causes of this trend were investigated by measuring the expression levels of three genes encoding upstream regulators of DNA repair protein expression, *BRCA1*, *E2F1* and *NRF2*. Expression of *BRCA1* and *NRF2* was decreased at 24 h of exposure (supplementary Table 2, available at Mutagenesis Online).

### Overall DNA methylation and specific methylation of DNA repair gene promoters

Since epigenetic modification, in particular DNA methylation, is well known to regulate gene expression (34), the overall methylation level of DNA was monitored, by 5-Me-dC quantification in A549 cells exposed for 4 h, 24 h or 48 h to TiO<sub>2</sub>-NPs (Table 2). No change in the overall DNA methylation level was detected by this assay. We therefore used RT-qPCR (EpiTect assay) to examine the methylation profile of 20 DNA repair gene promoters in cells exposed to TiO<sub>2</sub>-NPs. The promoters tested were those of *APE1*, *ATM*, *BRCA1*, *LIG3*, *MLH1*, *MRE11*, *MSH2*, *PMS2*, *POLB*, *RAD50*, *UNG*, *XPC* and *XRCC1*. Exposure of A549 cells to 100 µg/mL of TiO<sub>2</sub>-NPs for 4 or 24 h led to a moderate, but statistically significant, increase in methylation levels of *APE1* (at 4 and 24 h), *POLB* (at 4 h), *MRE11A* (at 24 h) and *PMS2* (at 24 h) (Figure 6).

### Discussion


This article compares the impact of TiO<sub>2</sub>-NPs on human bronchial BEAS-2B cells and alveolar A549 cells in terms of cell mortality,



**Figure 3.** Heat map representation of DNA excision/synthesis repair activities in A549 and BEAS-2B cells exposed to  $\text{TiO}_2$ -NPs. Cells were exposed to 100  $\mu\text{g}/\text{mL}$   $\text{TiO}_2$ -NPs for 4, 24 or 48 h (BEAS-2B); 24 or 48 h (A549). NT: non-treated cells (i.e. control cells); T: treated cells; 24 and 48 for 24 h and 48 h, respectively. Colour scale ranges between blue (lower value) and red (higher value). Each value represents the relative contribution of a specific repair pathway to total DNA repair activity in the cell after normalisation of data. For instance, when considering DNA repair activity in a chosen condition (NT or T, 24 or 48), the pathway with the reddest colour is the most active, while the bluest pathway is the least active. The experiment was performed once with three independent replicates.

genotoxicity, DNA repair activity and expression of DNA repair genes. It is important to note that the same procedure was used to expose both cell lines—the same exposure medium, duration and  $\text{TiO}_2$ -NP concentration. Thus, both cell lines received the same dose of NPs, initially coated with the same corona.  $\text{TiO}_2$ -NPs were observed to be moderately toxic to cells, with equivalent mortality rates in A549 cells and BEAS-2B cells. Oxidative damage to DNA was also observed in both cell lines, with DNA strand breaks

and/or alkali-labile sites as well as oxidative damage, producing 8-oxo-dGuo. Similar levels of these types of lesion were observed in A549 and BEAS-2B, with lesions appearing as soon as 4 h after exposure in both cell lines. However, strand breaks and/or alkali-labile sites tended to appear earlier in A549 than in BEAS-2B cells. With increased exposure durations, the frequency of this damage decreased slightly in A549 cells, in line with the increase in DNA repair activity previously observed (12). In BEAS-2B, extended



**Figure 4.** DNA lesion excision activity in A549 cells exposed to TiO<sub>2</sub>-NPs. A549 cells were exposed to 100 µg/mL TiO<sub>2</sub>-NPs for 24 or 48 h. *In vitro* excision activity was assessed using the GlycoSPOT assay. The fluorescence of each spot was normalised for the total fluorescence measured on the slide to determine the relative contribution of each repair pathway. This experiment was performed once with three independent replicates ( $n = 3$ ). Statistical significance is indicated as \* $P < 0.05$  for exposed vs. control.

**Table 1.** Comparison of the modulation of expression of proteins involved in NER and BER in A549 and BEAS-2B cells exposed to TiO<sub>2</sub>-NPs<sup>a</sup>


	4 h		24 h		48 h	
	A549	BEAS-2B	A549	BEAS-2B	A549	BEAS-2B
OGG1						
APE1			0.48 ± 0.16			
PARP1			0.72 ± 0.09		0.55 ± 0.11	
POLB		<b>0.62 ± 0.10</b>	<b>0.56 ± 0.06</b>	<b>0.67 ± 0.15</b>		
XRCC1	0.81 ± 0.07					
PCNA	<b>0.64 ± 0.08</b>	<b>0.76 ± 0.14</b>	<b>0.54 ± 0.08</b>	<b>0.58 ± 0.06</b>		
LIG3	0.75 ± 0.11		0.62 ± 0.12			
POLE						
LIG1						
hHR23B			<b>0.70 ± 0.07</b>			<b>0.57 ± 0.16</b>
XPC	0.54 ± 0.05		0.48 ± 0.06			
DDB1	0.66 ± 0.18		0.59 ± 0.15			
DDB2	0.66 ± 0.09		0.46 ± 0.13			
CSA			0.59 ± 0.08			
CSB	0.49 ± 0.14		0.47 ± 0.13			

<sup>a</sup>Expression levels for genes encoding DNA repair proteins were analysed by RT-qPCR in A549 and BEAS-2B cells exposed to 100 µg/mL TiO<sub>2</sub>-NPs and compared with the levels in unexposed (control) cells. Reference genes: *GAPDH*, *S18* and *CycloB*. Fold change is indicated only when expression was significantly different in exposed cells vs. unexposed cells. Boldface indicates that expression changed in both A549 and BEAS-2B cells. Statistical significance was assessed using REST 2009 software (28).

exposure duration increased the number of strand breaks and/or alkali-labile sites slightly, a result that is compatible with the decreased DNA repair ability observed in these cells. Our results are thus in agreement with the vast majority of studies published to date. Indeed, most previous studies also reported only moderate cell mortality (4), which often reaches a plateau not exceeding 30% (9,12,35–38). In addition, most genotoxicity studies performed on BEAS-2B or A549 cells exposed to comparable TiO<sub>2</sub>-NP concentrations also reported a TiO<sub>2</sub>-NP-induced increase in numbers of DNA breaks and/or DNA oxidation sites (12,20,39–47). Moreover, except in A549, at the earlier time point (24 h), the results presented here indicate that DNA repair activity is reduced in both cell lines upon exposure to TiO<sub>2</sub>-NPs. Delving deeper into the cellular response to TiO<sub>2</sub>-NP exposure, we observed that the relative importance of NER decreased considerably, while the relative decrease of the BER pathway was less extensive. This suggests that BEAS-2B and A549 cells preserve BER, which is necessary to repair oxidative damage caused by TiO<sub>2</sub>-NPs. This phenomenon is not linked to transcriptional regulation, since at these time points, very few DNA repair genes are

modulated. Based on these endpoints for assessing NP toxicity, we conclude that the response of the two cell lines is equivalent. A similar response of A549 and BEAS-2B cells following NP exposure was previously reported by Kain *et al.* (48); in their case, the NPs tested were manganese dioxide, silver (Ag), cerium oxide, cobalt oxide, ferrous oxide, nickel oxide and silicon dioxide (SiO<sub>2</sub>) (48). Thus, either cell line is appropriate for use as a biological model for nanotoxicity studies investigating cytotoxicity and genotoxicity.

The most striking difference between the two cell lines is the early response observed in A549 cells but not in BEAS-2B. At 4 h and 24 h of exposure to TiO<sub>2</sub>-NPs, a slight increase in DNA repair activities was observed in A549, this decrease correlated with generalised, intense downregulation of DNA repair genes. A549 cells thus respond to TiO<sub>2</sub>-NPs in a biphasic way, with (i) a first phase of intense deregulation of gene expression and (ii) a second phase that can be interpreted as an adaptive response, during which cells amplify their response towards oxidatively generated DNA lesions. In the first phase, 31 DNA repair genes or upstream regulators of the 44 are downregulated in at least one TiO<sub>2</sub>-NP exposure condition



**Figure 5.** Schematic representation of DNA repair pathways and modulation of the gene expression profile for the proteins involved in each pathway. Protein name is indicated in blue when expression of its gene is downregulated, in black when expression of its gene is unchanged, in red when expression of its gene is upregulated.

(exposure duration or NP concentration). Strikingly, these genes are all downregulated. This unexpected profile could be an indication that TiO<sub>2</sub>-NPs impact transcription overall in a non-specific way. However, unpublished transcriptomic analyses conducted in our laboratory refute this hypothesis; in these experiments, exposure of A549 cells to the same TiO<sub>2</sub>-NPs as in the present study induced an

overall upregulation of gene expression. A similar generalised downregulation of DNA repair gene expression was reported in lung cells exposed to Ag-NPs,(49) and in liver and lymphoma cells exposed to Ag-NPs (50,51). In these other cell types, expression of *OGG1*, *NRF2*, *XPA* and *ERCC2* was decreased (50,51). This inhibition may be related to the fact that Ag-NPs are thought to dissolve in the

intracellular compartment and to the affinity of Ag ions for thiol (-SH) groups. Some DNA repair proteins and transcription factors are zinc (Zn) finger proteins, in which Zn is bonded to the protein via -SH groups, the hypothesis is that Ag ions may replace Zn in the active site of these proteins, causing their inactivation (48). In contrast, TiO<sub>2</sub>-NPs are considered to be very chemically inert (e.g. they do not dissolve) they would not therefore engage in the same type of interactions as Ag-NPs and consequently must act via a different mechanism. Recently, DNA repair gene expression profiles in various species exposed to a range of nanomaterials showed that TiO<sub>2</sub>-NPs induce downregulation of five of the six DNA repair genes at 24 h of exposure (20). This effect may be produced by sequestration of transcription factors in the protein corona that forms on the surface of NPs. However, the fact that downregulation of gene expression is restricted to DNA repair genes suggests that one or more upstream regulator(s) of DNA repair processes may be directly affected by TiO<sub>2</sub>-NP exposure. Attempts to identify this (these) upstream regulator(s) led us to analyse the expression of *BRCA1*, *E2F1* and *NRF2* in TiO<sub>2</sub>-NP-exposed cells, as well as the overall methylation levels, and more specifically the methylation profile

**Table 2.** Overall DNA methylation level in A549 cells exposed to TiO<sub>2</sub>-NPs<sup>a</sup>


	4 h	24 h	48 h
Ctrl	0.95 ± 0.70	1.00 ± 0.55	0.80 ± 0.30
10 µg/mL TiO <sub>2</sub> -NPs	1.02 ± 0.57	0.98 ± 0.55	0.91 ± 0.05
100 µg/mL TiO <sub>2</sub> -NPs	1.04 ± 0.68	0.88 ± 0.34	0.92 ± 0.18

<sup>a</sup>Overall DNA methylation was assessed by measuring 5-methyl deoxycytosine in A549 cells exposed for 4, 24 or 48 h to 0, 10 or 100 µg/mL TiO<sub>2</sub>-NPs. Results show DNA methylation rate, average ± SD, *n* = 3.

of DNA repair gene promoters. Contrary to what was observed by Li *et al.* (52) in A549 cells exposed to carbon-based nanomaterials, we did not detect any overall change in DNA methylation levels. Moreover, when examining the promoter methylation profile, although some promoters were found to be more methylated in TiO<sub>2</sub>-NP-exposed cells than in control cells, the methylation levels observed in this study remained low compared with the levels reported in cancer cells (53). Only 4 of the 20 gene promoters tested here showed increased methylation levels. This suggests that methylation is unlikely to be the only factor behind the overall downregulation of DNA repair genes observed. Nevertheless, the *APE1* promoter shows increased methylation in NP-exposed cells, which is consistent with its decreased expression. While epigenetic alterations are known to be a major causative factor in diseases or adverse effects of environmental toxicants, especially those found in ambient air and due to ultrafine particles (54), based on our results, we believe that it is not a key mode of action of TiO<sub>2</sub>-NPs.

Bai *et al.* (55) reported altered methylation of the *PARP1* gene promoter in TiO<sub>2</sub>-NP-exposed cells, which correlated with oxidative stress (55). Although we did not assess methylation of the *PARP1* promoter in the present study, we did observe downregulation of this gene. Hypermethylation of *PARP1* promoter was also reported in HaCaT cells exposed to SiO<sub>2</sub>-NPs, where it was correlated with *PARP1* gene repression (56). Our data thus confirm the implication of epigenetic regulation, in particular methylation of gene promoters, in regulating the gene expression associated with TiO<sub>2</sub>-NP exposure, although it may not be the only causative factor.

Decreased expression of *NRF2* and *BRCA1* could also partly explain the reduced DNA repair activity observed here. Indeed, inactivation of *NRF2*, in addition to its antioxidant activities, has been shown to significantly slow down DNA repair processes (57). It especially affects HR as it binds to the antioxidant-response-element



**Figure 6.** DNA methylation profile of DNA repair gene promoters. The methylation level of each promoter was measured using the EpiTect® assay (Qiagen), on A549 cells exposed for 4 h (A, B) or 24 h (C, D) to 0 (Ctrl) or 100 µg/mL of TiO<sub>2</sub>-NPs. Statistical significance is indicated as \**P* < 0.05 for exposed vs. control.

sequences of some DNA repair proteins involved in this pathway (58). Similarly, breast cancer type 1 susceptibility protein (BRCA1) is known to be involved in the HR, BER and NER pathways (59), as it regulates transcription and has E3 ubiquitin ligase activity. Mutation of *BRCA1* compromises the repair of oxidatively damaged DNA and makes cells more sensitive to the alkylating agent methane methylsulphonate (59). We also reported similar effects in A549 cells following chronic exposure to TiO<sub>2</sub>-NPs (18).

The genes downregulated encode proteins from all the DNA repair pathways and from every stage of each pathway from damage recognition to DNA resynthesis and ligation. However, the excision stage appears to be less affected than others, as only one glycosylase (UNG) was downregulated of the three tested. This reduced sensitivity was confirmed by a multiplexed array of excision activities of A549 cells, which showed that only excision of A paired with 8-oxo-dGuo and EthA were affected in TiO<sub>2</sub>-NP-exposed cells. Contrary to expectations, OGG1 activity is not impaired in TiO<sub>2</sub>-NP-exposed cells, even though oxidative stress has been described as a potent and reversible inactivator of this protein (33). However, elimination of A paired with 8-oxo-dGuo, which occurs when 8-oxo-dGuo is bypassed by DNA polymerases during replication—causing G-to-T transversion mutations—is impaired in TiO<sub>2</sub>-NP-exposed cells. These lesions are primarily excised by MUTYH glycosylase, which is considered as a backup system for the removal of 8-oxo-dGuo (60). Inactivation of this repair pathway raises concerns as mutation of the *MUTYH* gene has been described in a common form of familial colorectal cancer (60).

Downregulation of the expression of this DNA repair gene, together with impaired DNA repair activity, proves that high concentrations of TiO<sub>2</sub>-NPs cause intense indirect genotoxicity, while lower concentrations do not. The TiO<sub>2</sub>-NP exposure doses used here were 0, 1, 10 or 100 µg/mL, which correspond to 0.17–17 µg/cm<sup>2</sup> or 1.1 pg/cell–0.11 ng/cell. The lower end of this dose range roughly corresponds to the alveolar deposition theoretically obtained after 24 h of exposure to a 1 mg/m<sup>3</sup> TiO<sub>2</sub>-NP aerosol (61). This equivalent aerosol concentration is between the maximum exposure concentrations recommended by the National Institute for Occupational Safety and Health (NIOSH) and the maximum recommended by New Energy and Industrial Technology Development Organization (NEDO) for TiO<sub>2</sub>-NPs, which are 0.3 and 1.2 mg/m<sup>3</sup>, respectively. The highest end of the dose range tested here could represent a worst-case scenario, but it is nevertheless a very high exposure concentration which could be considered unrealistic. However, occupational exposure is often chronic in nature, and we previously reported that chronicity increases the impact of TiO<sub>2</sub>-NPs as it is correlated with increased intracellular accumulation (17,18). Consequently, even if the high concentration is unrealistic as an acute dose, we consider that the acute exposure model used in the present study could nevertheless be informative from a risk assessment perspective.

To summarise, in this article, we show that A549 and BEAS-2B cell lines display similar responses to TiO<sub>2</sub>-NPs, characterised by moderate mortality, significant oxidatively generated DNA damage and impaired DNA repair activities. Although several DNA repair activities are dramatically affected, BER activities are somehow preserved, suggesting attempts by the cells to repair oxidatively generated DNA damage. Before the onset of this adaptive response, a first phase is observed in A549 cells, during which DNA repair genes are intensely deregulated. This transcriptional modulation mainly consists of downregulation affecting almost every stage of the different DNA repair pathways. Attempts to identify upstream causative factor(s) led to the conclusion that gene promoter methylation

as well as NRF2 and BRCA1 might be involved in the effects of TiO<sub>2</sub>-NP exposure on DNA repair. Thus, our results describe indirect genotoxicity of TiO<sub>2</sub>-NPs in two lung cell lines due to impaired DNA repair.

## Supplementary data

Supplementary Tables 1 and 2 and Supplementary Figure 1 are available at *Mutagenesis* Online.

## Funding

This work was supported by the Atomic Energy and Alternative Energies Commission through the ‘Toxicology’ research programme; and the European Commission’s 7th Framework Programme project NanoMILE (contract no. NMP4-LA-2013-310451). It is a contribution to the Labex Serenade (n° ANR-11-LABX-0064) funded by the French Government’s ‘Investissements d’Avenir’ programme managed by the French National Research Agency (ANR) through the A\*MIDEX project (n° ANR-11-IDEX-0001-02).

Conflict of interest statement: None declared.

## References

- Shi, H., Magaye, R., Castranova, V. and Zhao, J. (2013) Titanium dioxide nanoparticles: a review of current toxicological data. *Part. Fibre Toxicol.*, 10, 15.
- Boland, S., Hussain, S. and Baeza-Squiban, A. (2014) Carbon black and titanium dioxide nanoparticles induce distinct molecular mechanisms of toxicity. *Nanomed. Nanobiotechnol.*, 6, 641–652.
- Johnston, H. J., Hutchison, G. R., Christensen, F. M., Peters, S., Hankin, S. and Stone, V. (2009) Identification of the mechanisms that drive the toxicity of TiO<sub>2</sub> particulates: the contribution of physicochemical characteristics. *Part. Fibre Toxicol.*, 6, 33.
- Landsiedel, R., Sauer, U. G., Ma-Hock, L., Schneckeburger, J. and Wiemann, M. (2014) Pulmonary toxicity of nanomaterials: a critical comparison of published in vitro assays and in vivo inhalation or instillation studies. *Nanomedicine (Lond.)*, 9, 2557–2585.
- Skocaj, M., Filipic, M., Petkovic, J. and Novak, S. (2011) Titanium dioxide in our everyday life; is it safe? *Radiol. Oncol.*, 45, 227–247.
- Baan, R. A. (2007) Carcinogenic hazards from inhaled carbon black, titanium dioxide, and talc not containing asbestos or asbestiform fibers: recent evaluations by an IARC Monographs Working Group. *Inhal. Toxicol.*, 19(Suppl 1), 213–228.
- Gonzalez, L. and Kirsch-Volders, M. (2016) Biomonitoring of genotoxic effects for human exposure to nanomaterials: the challenge ahead. *Mutat. Res.*, 768, 14–26.
- Bruinink, A., Wang, J. and Wick, P. (2015) Effect of particle agglomeration in nanotoxicology. *Arch. Toxicol.*, 89, 659–675.
- Simon-Deckers, A., Gouget, B., Mayne-L’hermite, M., Herlin-Boime, N., Reynaud, C. and Carrière, M. (2008) In vitro investigation of oxide nanoparticle and carbon nanotube toxicity and intracellular accumulation in A549 human pneumocytes. *Toxicology*, 253, 137–146.
- Ahlinder, L., Ekstrand-Hammarstrom, B., Geladi, P. and Osterlund, L. (2013) Large uptake of titania and iron oxide nanoparticles in the nucleus of lung epithelial cells as measured by Raman imaging and multivariate classification. *Biophys. J.*, 105, 310–319.
- Geiser, M., Rothen-Rutishauser, B., Kapp, N., et al. (2005) Ultrafine particles cross cellular membranes by nonphagocytic mechanisms in lungs and in cultured cells. *Environ. Health Perspect.*, 113, 1555–1560.
- Jugan, M. L., Barillet, S., Simon-Deckers, A., Herlin-Boime, N., Sauvaigo, S., Douki, T. and Carrière, M. (2012) Titanium dioxide nanoparticles exhibit genotoxicity and impair DNA repair activity in A549 cells. *Nanotoxicology*, 6, 501–513.

13. Li, N., Ma, L., Wang, J., *et al.* (2009) Interaction between nano-anatase TiO<sub>2</sub> and liver DNA from mice *in vivo*. *Nanoscale Res. Lett.*, 5, 108–115.
14. Shukla, R. K., Sharma, V., Pandey, A. K., Singh, S., Sultana, S. and Dhawan, A. (2011) ROS-mediated genotoxicity induced by titanium dioxide nanoparticles in human epidermal cells. *Toxicol. In Vitro*, 25, 231–241.
15. Magdolenova, Z., Collins, A., Kumar, A., Dhawan, A., Stone, V. and Dusinska, M. (2014) Mechanisms of genotoxicity. A review of *in vitro* and *in vivo* studies with engineered nanoparticles. *Nanotoxicology*, 8, 233–278.
16. Chen, T., Yan, J. and Li, Y. (2014) Genotoxicity of titanium dioxide nanoparticles. *J Food Drug Anal.*, 22, 95–104.
17. Armand, L., Biola-Clier, M., Bobyk, L., *et al.* (2016) Molecular responses of alveolar epithelial A549 cells to chronic exposure to titanium dioxide nanoparticles: A proteomic view. *J. Proteomics*, 134, 163–173.
18. Armand, L., Tarantini, A., Beal, D., *et al.* (2016) Long-term exposure of A549 cells to titanium dioxide nanoparticles induces DNA damage and sensitizes cells towards genotoxic agents. *Nanotoxicology*, 10, 913–923.
19. Huang, S., Chueh, P. J., Lin, Y. W., Shih, T. S. and Chuang, S. M. (2009) Disturbed mitotic progression and genome segregation are involved in cell transformation mediated by nano-TiO<sub>2</sub> long-term exposure. *Toxicol. Appl. Pharmacol.*, 241, 182–194.
20. Lan, J., Gou, N., Gao, C., He, M. and Gu, A.Z. (2014) Comparative and mechanistic genotoxicity assessment of nanomaterials via a quantitative toxicogenomics approach across multiple species. *Environ. Sci. Technol.*, 48, 12937–12945.
21. Tilton, S. C., Karin, N. J., Tolic, A., *et al.* (2014) Three human cell types respond to multi-walled carbon nanotubes and titanium dioxide nanobelts with cell-specific transcriptomic and proteomic expression patterns. *Nanotoxicology*, 8, 533–548.
22. ICRP. (1994) Human respiratory tract model for radiological protection. A report of a Task Group of the International Commission on Radiological Protection. *Ann. ICRP*, 24, 1–482.
23. Singh, A., Misra, V., Thimmulappa, R. K., *et al.* (2006) Dysfunctional KEAP1-NRF2 interaction in non-small-cell lung cancer. *PLoS Med.*, 3, e420.
24. Brun, E., Barreau, F., Veronesi, G., *et al.* (2014) Titanium dioxide nanoparticle impact and translocation through *ex vivo*, *in vivo* and *in vitro* gut epithelia. *Part. Fibre Toxicol.*, 11, 1743–8977.
25. Tice, R. R., Agurell, E., Anderson, D., *et al.* (2000) Single cell gel/comet assay: guidelines for *in vitro* and *in vivo* genetic toxicology testing. *Environ. Mol. Mutagen.*, 35, 206–221.
26. Frelon, S., Douki, T., Ravanat, J. L., Pouget, J. P., Tornabene, C. and Cadet, J. (2000) High-performance liquid chromatography-tandem mass spectrometry measurement of radiation-induced base damage to isolated and cellular DNA. *Chem. Res. Toxicol.*, 13, 1002–1010.
27. Pfaffl, M. W., Tichopad, A., Prgomet, C. and Neuvians, T. P. (2004) Determination of stable housekeeping genes, differentially regulated target genes and sample integrity: BestKeeper–Excel-based tool using pair-wise correlations. *Biotechnol. Lett.*, 26, 509–515.
28. Pfaffl, M. W. (2001) A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res.*, 29, e45.
29. Millau, J. F., Raffin, A. L., Caillaud, S., *et al.* (2008) A microarray to measure repair of damaged plasmids by cell lysates. *Lab Chip*, 8, 1713–1722.
30. Pons, B., Belmont, A. S., Masson-Genteuil, G., Chapuis, V., Oddos, T. and Sauvaigo, S. (2010) Age-associated modifications of Base Excision Repair activities in human skin fibroblast extracts. *Mech. Ageing Dev.*, 131, 661–665.
31. Moller, P., Jensen, D. M., Christophersen, D. V., *et al.* (2015) Measurement of oxidative damage to DNA in nanomaterial exposed cells and animals. *Environ. Mol. Mutagen.*, 56, 97–110.
32. Hoeijmakers, J. H. (2001) Genome maintenance mechanisms for preventing cancer. *Nature*, 411, 366–374.
33. Bravard, A., Vacher, N., Gouget, B., Coutant, A., de Boisferon, F. H., Marsin, S., Chevillard, S. and Radicella, J. P. (2006) Redox regulation of human OGG1 activity in response to cellular oxidative stress. *Mol. Cell Biol.*, 26, 7430–7436.
34. Siegfried, Z. and Simon, I. (2010) DNA methylation and gene expression. *Systems Biol. Med.*, 2, 362–371.
35. Chusuei, C. C., Wu, C. H., Mallavarapu, S., Hou, F. Y., Hsu, C. b.M., Winiarz, J. G., Aronstam, R. S. and Huang, Y. W. (2013) Cytotoxicity in the age of nano: the role of fourth period transition metal oxide nanoparticle physicochemical properties. *Chem. Biol. Interact.*, 206, 319–326.
36. Ghosh, M., Chakraborty, A. and Mukherjee, A. (2013) Cytotoxic, genotoxic and the hemolytic effect of titanium dioxide (TiO<sub>2</sub>) nanoparticles on human erythrocyte and lymphocyte cells *in vitro*. *J Appl. Toxicol.*, 33, 1097–1110.
37. Pujalte, I., Passagne, I., Brouillaud, B., Treguer, M., Durand, E., Ohayon-Courtes, C. and L'Azou, B. (2011) Cytotoxicity and oxidative stress induced by different metallic nanoparticles on human kidney cells. *Part. Fibre Toxicol.*, 8, 10.
38. Wang, J. J., Sanderson, B. J. and Wang, H. (2007) Cyto- and genotoxicity of ultrafine TiO<sub>2</sub> particles in cultured human lymphoblastoid cells. *Mutat. Res.*, 628, 99–106.
39. Falck, G. C., Lindberg, H. K., Suhonen, S., Vippola, M., Vanhala, E., Catalan, J., Savolainen, K. and Norppa, H. (2009) Genotoxic effects of nano-sized and fine TiO<sub>2</sub>. *Human Exp. Toxicol.*, 28, 339–352.
40. Gurr, J. R., Wang, A. S., Chen, C. H. and Jan, K. Y. (2005) Ultrafine titanium dioxide particles in the absence of photoactivation can induce oxidative damage to human bronchial epithelial cells. *Toxicology*, 213, 66–73.
41. Kansara, K., Patel, P., Shah, D., Shukla, R. K., Singh, S., Kumar, A. and Dhawan, A. (2015) TiO<sub>2</sub> nanoparticles induce DNA double strand breaks and cell cycle arrest in human alveolar cells. *Environ. Mol. Mutagen.*, 56, 204–217.
42. Karlsson, H. L., Cronholm, P., Gustafsson, J. and Moller, L. (2008) Copper oxide nanoparticles are highly toxic: a comparison between metal oxide nanoparticles and carbon nanotubes. *Chem. Res. Toxicol.*, 21, 1726–1732.
43. Karlsson, H. L., Gustafsson, J., Cronholm, P. and Möller, L. (2009) Size-dependent toxicity of metal oxide particles—a comparison between nano- and micrometer size. *Toxicol. Lett.*, 188, 112–118.
44. Prasad, R. Y., Wallace, K., Daniel, K. M., *et al.* (2013) Effect of treatment media on the agglomeration of titanium dioxide nanoparticles: impact on genotoxicity, cellular interaction, and cell cycle. *ACS Nano*, 7, 1929–1942.
45. Srivastava, R. K., Rahman, Q., Kashyap, M. P., Lohani, M. and Pant, A. B. (2011) Ameliorative effects of dimethylthiourea and N-acetylcysteine on nanoparticles induced cyto-genotoxicity in human lung cancer cells-A549. *PLoS One*, 6, e25767.
46. Srivastava, R. K., Rahman, Q., Kashyap, M. P., Singh, A. K., Jain, G., Jahan, S., Lohani, M., Lantow, M. and Pant, A. B. (2013) Nano-titanium dioxide induces genotoxicity and apoptosis in human lung cancer cell line, A549. *Human Exp. Toxicol.*, 32, 153–166.
47. Wang, Y., Cui, H., Zhou, J., Li, F., Wang, J., Chen, M. and Liu, Q. (2015) Cytotoxicity, DNA damage, and apoptosis induced by titanium dioxide nanoparticles in human non-small cell lung cancer A549 cells. *Environ. Sci. Pollut. Res.*, 22, 5519–5530.
48. Kain, J., Karlsson, H. L. and Möller, L. (2012) DNA damage induced by micro- and nanoparticles—interaction with FPG influences the detection of DNA oxidation in the comet assay. *Mutagenesis*, 27, 491–500.
49. Asharani, P., Sethu, S., Lim, H. K., Balaji, G., Valiyaveetil, S. and Hande, M. P. (2012) Differential regulation of intracellular factors mediating cell cycle, DNA repair and inflammation following exposure to silver nanoparticles in human cells. *Genome Integr.*, 3, 2.
50. Mei, N., Zhang, Y., Chen, Y., Guo, X., Ding, W., Ali, S. F., Biris, A. S., Rice, P., Moore, M. M. and Chen, T. (2012) Silver nanoparticle-induced mutations and oxidative stress in mouse lymphoma cells. *Environ. Mol. Mutagen.*, 53, 409–419.
51. Piao, M. J., Kim, K. C., Choi, J. Y., Choi, J. and Hyun, J. W. (2011) Silver nanoparticles down-regulate Nrf2-mediated 8-oxoguanine DNA glycosylase 1 through inactivation of extracellular regulated kinase and protein kinase B in human Chang liver cells. *Toxicol. Lett.*, 207, 143–148.
52. Li, J., Tian, M., Cui, L., Dwyer, J., Fullwood, N. J., Shen, H. and Martin, F. L. (2016) Low-dose carbon-based nanoparticle-induced effects in A549 lung cells determined by biospectroscopy are associated with increases in genomic methylation. *Scientific Rep.*, 6, 20207.

53. Lahtz, C. and Pfeifer, G. P. (2011) Epigenetic changes of DNA repair genes in cancer. *J. Mol. Cell Biol.*, 3, 51–58.
54. Stocco, A., Karlsson, H. L., Coppède, F. and Migliore, L. (2013) Epigenetic effects of nano-sized materials. *Toxicology*, 313, 3–14.
55. Bai, W., Chen, Y. and Gao, A. (2015) Cross talk between poly(ADP-ribose) polymerase 1 methylation and oxidative stress involved in the toxic effect of anatase titanium dioxide nanoparticles. *Int. J. Nanomed.*, 10, 5561–5569.
56. Gong, C., Tao, G., Yang, L., Liu, J., Liu, Q., Li, W. and Zhuang, Z. (2012) Methylation of PARP-1 promoter involved in the regulation of nano-SiO<sub>2</sub>-induced decrease of PARP-1 mRNA expression. *Toxicol. Lett.*, 209, 264–269.
57. Huang, Y., Li, W., Su, Z. Y. and Kong, A. N. (2015) The complexity of the Nrf2 pathway: beyond the antioxidant response. *J. Nutr. Biochem.*, 26, 1401–1413.
58. Jayakumar, S., Pal, D. and Sandur, S. K. (2015) Nrf2 facilitates repair of radiation induced DNA damage through homologous recombination repair pathway in a ROS independent manner in cancer cells. *Mutat. Res.*, 779, 33–45.
59. Alli, E. and Ford, J. M. (2015) BRCA1: Beyond double-strand break repair. *DNA Repair*, 32, 165–171.
60. David, S. S., O'Shea, V. L. and Kundu, S. (2007) Base-excision repair of oxidative DNA damage. *Nature*, 447, 941–950.
61. Gangwal, S., Brown, J. S., Wang, A., Houck, K. A., Dix, D. J., Kavlock, R. J. and Hubal, E. A. (2011) Informing selection of nanomaterial concentrations for ToxCast in vitro testing based on occupational exposure potential. *Environ. Health Perspect.*, 119, 1539–1546.